
Action Steps for Students, Faculty, and Staff to Prevent the Spread of Flu

August 20, 2009 3:00 PM ET

The Centers for Disease Control and Prevention (CDC) recommends 4 main ways you may keep from getting sick with the flu:

1. **Practice good hand hygiene** by washing your hands often with soap and water, especially after coughing or sneezing. Alcohol-based hand cleaners are also effective.
2. **Cover your mouth and nose with a tissue when you cough or sneeze.**
If you don't have a tissue, cough or sneeze into your elbow or shoulder; not into your hands.
3. **Stay home or at your place of residence if you are sick** for at least 24 hours after you no longer have a fever (100 degrees Fahrenheit or 38 degrees Celsius) or signs of a fever (have chills, feel very warm, have a flushed appearance, or are sweating). This should be determined without the use of fever-reducing medications (any medicine that contains ibuprofen or acetaminophen). Staying away from others while sick can prevent others from getting sick too. Ask a roommate, friend, or family member to check up on you and to bring you food and supplies if needed.
4. **Talk to your health care provider** to find out if you should be vaccinated for seasonal flu and/or 2009 H1N1 flu. Information about 2009 H1N1 flu vaccination can be found at: www.cdc.gov/h1n1flu/vaccination. Information about seasonal flu vaccine can be found at: www.cdc.gov/flu/protect/keyfacts.htm.